

## RESEARCH ARTICLES

# Identifying the Basal Angiosperm Node in Chloroplast Genome Phylogenies: Sampling One's Way Out of the Felsenstein Zone

Jim Leebens-Mack,\* Linda A. Raubeson,† Liying Cui,\* Jennifer V. Kuehl,‡ Matthew H. Fourcade,‡ Timothy W. Chumley,§ Jeffrey L. Boore,‡|| Robert K. Jansen,§ and Claude W. dePamphilis\*

\*Department of Biology, Institute of Molecular Evolutionary Genetics, and The Huck Institutes of Life Sciences, The Pennsylvania State University; †Department of Biological Sciences, Central Washington University; ‡Department of Energy Joint Genome Institute and University of California Lawrence Berkeley National Laboratory, Walnut Creek, California; §Section of Integrative Biology and Institute of Cellular and Molecular Biology, University of Texas, Austin; and ||Department of Integrative Biology, University of California, Berkeley

While there has been strong support for *Amborella* and Nymphaeales (water lilies) as branching from basal-most nodes in the angiosperm phylogeny, this hypothesis has recently been challenged by phylogenetic analyses of 61 protein-coding genes extracted from the chloroplast genome sequences of *Amborella*, *Nymphaea*, and 12 other available land plant chloroplast genomes. These character-rich analyses placed the monocots, represented by three grasses (Poaceae), as sister to all other extant angiosperm lineages. We have extracted protein-coding regions from draft sequences for six additional chloroplast genomes to test whether this surprising result could be an artifact of long-branch attraction due to limited taxon sampling. The added taxa include three monocots (*Acorus*, *Yucca*, and *Typha*), a water lily (*Nuphar*), a ranunculid (*Ranunculus*), and a gymnosperm (*Ginkgo*). Phylogenetic analyses of the expanded DNA and protein data sets together with microstructural characters (indels) provided unambiguous support for *Amborella* and the Nymphaeales as branching from the basal-most nodes in the angiosperm phylogeny. However, their relative positions proved to be dependent on the method of analysis, with parsimony favoring *Amborella* as sister to all other angiosperms and maximum likelihood (ML) and neighbor-joining methods favoring an *Amborella* + Nymphaeales clade as sister. The ML phylogeny supported the later hypothesis, but the likelihood for the former hypothesis was not significantly different. Parametric bootstrap analysis, single-gene phylogenies, estimated divergence dates, and conflicting indel characters all help to illuminate the nature of the conflict in resolution of the most basal nodes in the angiosperm phylogeny. Molecular dating analyses provided median age estimates of 161 MYA for the most recent common ancestor (MRCA) of all extant angiosperms and 145 MYA for the MRCA of monocots, magnoliids, and eudicots. Whereas long sequences reduce variance in branch lengths and molecular dating estimates, the impact of improved taxon sampling on the rooting of the angiosperm phylogeny together with the results of parametric bootstrap analyses demonstrate how long-branch attraction might mislead genome-scale phylogenetic analyses.

## Introduction

Characterized by Darwin (1903) as “an abominable mystery,” the early radiation of angiosperms was pivotal in the evolutionary history of our biota. After years of controversy concerning identity of the basal-most node in the angiosperm phylogeny, a series of studies using multiple genes from the chloroplast, mitochondrial, and nuclear genomes identified *Amborella*, the Nymphaeales, and Austrobaileyales as successive sister lineages relative to all other angiosperms (Mathews and Donoghue 1999; Parkinson, Adams, and Palmer 1999; Qiu et al. 1999; P. S. Soltis, D. E. Soltis, and Chase 1999; Barkman et al. 2000; Graham and Olmstead 2000; Zanis et al. 2002; Borsch et al. 2003; Hilu et al. 2003; Stefanović, Rice, and Palmer 2004; e.g., fig. 1A). Many of these studies found that the inferred relationship between *Amborella* and the Nymphaeales varied when using differing methods of phylogenetic reconstruction, models of molecular evolution, and subsets of taxa (Barkman et al. 2000; Graham and Olmstead 2000; Zanis et al. 2002; Stefanović, Rice, and Palmer 2004), with each lineage sometimes inferred to be most basal and, in some cases, for the two to form a single

clade, sister to all other angiosperm lineages (e.g., fig. 1B and C). While the branching order of *Amborella* and the Nymphaeales relative to each other and the rest of the angiosperms has remained controversial, there has been widespread consensus in the recent plant systematics literature for *Amborella* and Nymphaeales branching off at the base of the angiosperm phylogeny, followed subsequently by Austrobaileyales and the remaining angiosperm lineages.

That consensus was recently challenged by the results of phylogenetic analyses of 61 protein-coding genes common to 14 chloroplast genome sequences including the recently sequenced plastid genomes of *Amborella trichopoda* (Goremykin et al. 2003) and *Nymphaea alba* (Goremykin et al. 2004). The analyses of Goremykin and colleagues placed the monocots, represented by the chloroplast genomes of rice (*Oryza sativa*), maize (*Zea mays*), and wheat (*Triticum aestivum*), as sister to all other extant lineages in the angiosperm phylogeny. This result is quite intriguing because much of our current understanding of morphological, developmental, and molecular evolution in early angiosperm history would have to change if monocots are in fact sister to all other angiosperms (e.g., fig. 1D). As Goremykin et al. (2004, p. 1452) point out, however, the hypothesized basal divergence of monocots from all other angiosperms must be tested in analyses with increased taxon sampling. Genome-scale phylogenetic studies, where character sampling is very deep but taxon sampling is typically sparse, are particularly susceptible to long-branch

Key words: angiosperm phylogeny, phylogenomics, parametric bootstrap.

E-mail: jleebensmack@psu.edu.

*Mol. Biol. Evol.* 22(10):1948–1963. 2005

doi:10.1093/molbev/msi191

Advance Access publication June 8, 2005




FIG. 1.—Nucleotide-based ML phylogenies estimated using the HKY +  $\Gamma$  + I substitution model while constraining *Amborella* (A), *Amborella* + water lilies (B), water lilies (C), or monocots (D) as sister to remaining angiosperms. Likelihood for each hypothesis is shown with each phylogeny.

attraction (Soltis et al. 2004b; Philippe, Lartillot, and Brinkmann 2005), and multiple lines of evidence suggest that the placement of the grasses in the analyses of Goremykin et al. (2003, 2004) may be an artifact of sparse taxon sampling, particularly within the monocots (D. E. Soltis and P. S. Soltis 2004; Soltis et al. 2004b; Stefanović, Rice, and Palmer 2004; but see Lockhart and Penny 2005; Martin et al. 2005). Here we test this hypothesis directly by adding the corresponding 61-gene data sets for six additional species, including three nongrass monocot species, a ranunculid (the sister clade to all other eudicots), an additional water lily, and the gymnosperm *Ginkgo biloba*. Our study also includes an evaluation of microstructural mutations, a class of evolutionary events that may be less subject to homoplastic events than base substitutions. In addition, we extend the earlier work of Zanis et al. (2002) using parametric bootstrap analyses to investigate estimated branching events at the base of the angiosperms and estimate minimum divergence dates for well-supported clades on the phylogeny.

## Materials and Methods

### Sequencing

Templates suitable for constructing random insert plasmid libraries were generated for the chloroplast genomes of *G. biloba*, *Nuphar advena*, *Acorus americanus*, *Yucca schidigera*, *Typha latifolia*, and *Ranunculus macranthus* by one of the three ways (Jansen et al. 2005): (1) sucrose gradient isolation of pure chloroplast DNA (cpDNA) (*Ginkgo*, *Nuphar*, and *Ranunculus*), (2) rolling circle am-

plification of the entire plastid genome using plastid isolations (*Acorus* and *Typha*), and (3) cpDNA-containing clones identified through screening of a fosmid genomic library (*Yucca*). The low-coverage (<0.5 $\times$ ) *Yucca* fosmid library was produced using the Epicentre CopyControl<sup>TM</sup> Fosmid Library Production Kit according to manufacturer's protocols (<http://www.epibio.com>). All templates were sheared by serial passage through a narrow aperture using a Hydroshear<sup>®</sup> device (Genomic Solutions, Ann Arbor, Mich.) into random fragments of approximately 3 kb, and then cloned into the pUC18 plasmid vector to create a clone library. These clones were robotically processed through colony picking, rolling circle amplification using TempliPhi<sup>TM</sup> (Amersham Biosciences, Little Chalfont, UK) sequencing reactions with either ET terminators (Amersham Biosciences) or BigDye<sup>TM</sup> terminators Applied Biosystems, Foster City, Calif.), and then reads of approximately 700 nt each were determined from each end of each clone. Detailed protocols are available at [http://www.jgi.doe.gov/sequencing/protocols/protos\\_production.html](http://www.jgi.doe.gov/sequencing/protocols/protos_production.html). Sequence reads were trimmed and assembled using phred and phrap (Ewing and Green 1998; Ewing et al. 1998) and manually interpreted using Consed (Gordon, Abajian, and Green 1998) and Sequencher (<http://www.genecodes.com>). Roughly 4,000 sequencing reads were determined for each chloroplast genome, giving about 2.8 million nucleotides of sequence (at Q20 or greater quality), for an average depth of coverage in the assemblies of about 12 $\times$  (considering losses due to impure cpDNA preparations). The sequence of each cpDNA was finished to meet the quality criteria specified in Jansen et al. (2005) by additional

**Table 1**  
**GenBank Accession Numbers for the Sequences Included in This Study**

Taxon (Collection locale: voucher ID)	GenBank Accession Numbers	Reference
<b>Bryophytes</b>		
<i>Anthoceros formosae</i>	NC_004543	Kugita et al. 2003b
<i>Physcomitrella patens</i>	NC_005087	Sugiura et al. 2003
<i>Marchantia polymorpha</i>	NC_001319	Ohyama et al. 1988
<b>Ferns and allies</b>		
<i>Psilotum nudum</i>	NC_003386	Wakasugi et al., 1998
<i>Adiantum capillus-veneris</i>	NC_004766	Wolf et al. 2003
<b>Gymnosperms</b>		
<i>Pinus thunbergii</i>	NC_001631	Wakasugi et al. 1994
<i>Ginkgo biloba</i> (Travis Co., Tex.; RCH155 TEX)	DQ069337–DQ069702	Current study
<b>Basal-most angiosperm lineages<sup>a</sup></b>		
<i>Amborella trichopoda</i>	NC_005086	Goremykin et al. 2003
<i>Nymphaea alba</i>	NC_006050	Goremykin et al. 2004
<i>Nuphar advena</i> (Centre Co., Pa.; PAC 95537 PAC)	DQ069337–DQ069702	Current study
<b>Monocots</b>		
<i>Acorus americanus</i> (Crawford Co., Pa.; PAC 95538 PAC)	DQ069337–DQ069702	Current study
<i>Typha latifolia</i> (Yavapai Co., Ariz.; RCH188 TEX)	DQ069337–DQ069702	Current study
<i>Yucca schidigera</i> (San Diego Co., Calif.; jlm-yuc370 PAC)	DQ069337–DQ069702	Current study
<i>Saccharum officinarum</i>	NC_006084	Asano et al. 2004
<i>Zea mays</i>	NC_001666	Maier et al. 1995
<i>Oryza sativa</i>	NC_001320	Hiratsuka et al. 1989
<i>Triticum aestivum</i>	NC_002762	Ogihara et al. 2000
<b>Magnoliids</b>		
<i>Calycanthus floridus</i>	NC_004993	Goremykin et al. 2003
<b>Eudicots</b>		
<i>Ranunculus macranthus</i> (Travis Co., Tex.; RCH1184 TEX)	DQ069337–DQ069702	Current study
<i>Nicotiana tabacum</i>	NC_001879	Shinozaki et al. 1986
<i>Atropa belladonna</i>	NC_004561	Schmitz-Linneweber et al. 2002
<i>Spinacia oleracea</i>	NC_002202	Schmitz-Linneweber et al. 2001
<i>Lotus corniculatus</i>	NC_002694	Kato et al. 2000
<i>Medicago truncatula</i>	NC_003119	Lin S., H. Wu, H. Jia, P. Zhang, R. Dixon, G. May, R. Gonzales, and B. A. Roe, unpublished data
<i>Arabidopsis thaliana</i>	NC_000932	Sato et al. 1999
<i>Oenothera elata</i>	NC_002693	Hupfer et al. 2000

<sup>a</sup> Basal angiosperm lineages as determined in most molecular systematic studies since 1999 (see text).

sequencing reactions that targeted gaps or parts of the genomes with potential errors using primers specifically designed for targeted portions of the plastid genome.

### Phylogenetic Analyses

The 61 genes included in the analyses of Goremykin et al. (2003, 2004) were extracted from high-quality contigs (Q values >40 for all extracted bases) of our six new chloroplast genome sequences using the organellar genome annotation program DOGMA (<http://evogen.jgi-psf.org>; Wyman, Jansen, and Boore 2004). The same set of 61 genes was extracted from chloroplast genome sequences for the 18 other available species (table 1). Inferred amino acid sequences for each of the 61 genes were aligned using ClustalW (Thompson, Higgins, and Gibson 1994) and adjusted manually. A nucleotide alignment was then forced to correspond to the amino acid alignment and further adjusted.

The complete amino acid and nucleotide alignments are available in our Chloroplast Genome Database (<http://chloroplast.cbio.psu.edu/>), and sequences are available in GenBank (accession numbers DQ069337–DQ069702; table 1).

Phylogenetic analyses of nucleotide alignments using maximum parsimony (MP) and Neighbor-Joining (NJ) were performed using PAUP\* version 4.10 (Swofford 2003), and maximum likelihood (ML) analyses were performed using both PAUP\* and PHYML version 2.4.4 (Guindon and Gascuel 2003). All MP searches were heuristic with 10 random addition replicates and Tree Bisection-Reconnection branch swapping. The Hasegawa-Kishino-Yano (HKY) (Hasegawa, Kishino, and Yano 1985) model of molecular evolution was used in ML and NJ analyses of the nucleotide alignments. ML estimates of HKY distances were used for the NJ analyses. ML and NJ analyses of

amino acid alignments were performed using PHYML and PHYLIP version 3.63 (seqboot, protdist, and neighbor; Felsenstein 2004), respectively, under the Jones-Taylor-Thornton (JTT) model (Jones, Taylor, and Thornton 1992). ML and NJ analyses of the nucleotide alignment were run with and without rate variation among sites (HKY +  $\Gamma$ ), with and without invariant sites (HKY +  $\Gamma$  + I). ML and NJ analyses of the amino acid alignment were run with and without rate variation among sites (JTT +  $\Gamma$ ), and ML analyses were also run with and without invariant sites (JTT +  $\Gamma$  + I). Rate variation among sites was estimated as a discrete gamma distribution (Yang 1994) with six rate classes. ML parameter estimates for rate variation across sites and for invariant sites were optimized simultaneously with topology and branch lengths in PHYML. In order to avoid being trapped in local optima, all ML analyses performed in PHYML were run with five starting trees including the BIONJ tree (default) and the four trees shown in figure 1. Likelihood ratio tests showed significant improvement in the fits of both nucleotide and protein evolution models with the addition of parameters for rate variation across sites and invariant sites ( $P \ll 0.001$ ). Non-parametric bootstrap analyses (Felsenstein 1985) were performed for all analyses with 200 pseudoreplicates.

In addition to the typical ML analysis, constrained likelihood trees were estimated for the nucleotide alignment setting one of four lineages as sister to the remaining angiosperms: *Amborella* (fig. 1A), *Amborella* + Nymphaeales (fig. 1B), Nymphaeales (fig. 1C), and monocots (fig. 1D; Goremykin et al. 2003, 2004). The Shimodaira-Hasegawa (SH) (Shimodaira and Hasegawa 1999) test was performed in PAUP\* to determine whether any of the resulting phylogenies were significantly worse than the ML tree. The test was performed using 10,000 bootstrap replicates. MP topologies were tested similarly in PAUP\* using the Wilcoxon test.

In their analyses, Goremykin et al. (2003, 2004) removed third codon positions from their nucleotide alignments citing concerns over saturation of nucleotide substitutions at synonymous sites. Pairwise genetic distances for all taxon pairs in our analysis (HKY model) were calculated separately for codon positions 1 + 2 and position 3. A linear relationship was observed between genetic divergence (HKY distances) at third versus first and second positions (fig. 2), suggesting that saturation was not seriously biasing distance estimates, so third positions were included in our phylogenetic analyses. The inclusion of third codon positions changed bootstrap values for some nodes, but with the exception of the poorly supported placement of *Calycanthus*, the optimal MP, ML, and NJ topologies were the same in 61-gene analyses with and without their inclusion (see *Results*).

Initial analyses of codon usage, amino acid content, and branch lengths showed that the fern *Adiantum* and hornwort *Anthoceros* sequences are extremely divergent from their closest relatives in the study, in part due to extensive RNA editing (Kugita et al. 2003b; Wolf et al. 2003). These taxa, which were not critical to our study, were eliminated from further analyses. Gapped sites were also excluded from additional analyses because these often represented regions of questionable annotation and align-




Fig. 2.—Pairwise HKY distances estimated for third codon positions are linearly correlated with distances estimated for first and second positions.

ment. An exception was made for well-aligned sites in *rpoA* and *ccsA*, which are missing from the plastome of *Physcomitrella patens* and were coded as gap for this taxon. The resulting nucleotide and amino acid alignments included 39,978 and 13,326 characters, respectively. Exclusion of *Adiantum*, *Anthoceros*, or gapped sites affected only the poorly supported placement of *Calycanthus* relative to the eudicots in MP, NJ, and ML topologies (data not shown).

Ané et al. (2005) have recently shown among-lineage rate heterogeneity, or heterotachy (Lopez, Casane, and Philippe 2002), at many sites in plant plastid genes. This type of rate variation can confound phylogenetic analyses given some patterns of variation (e.g., Kolaczkowski and Thornton 2004; Spencer, Susko, and Roger 2005). We attempted to control for heterotachy in distance analyses of both nucleotide and amino acid alignments by estimating LogDet distances (Lake 1994; Lockhart et al. 1994; Steel 1994) with and without among-site rate heterogeneity using LDDist (Thollessen 2004; Martin et al. 2005).

Parsimony analyses were also performed on all 61 genes individually. Analyses were performed on alignments of all three codon positions, first and second codon positions and amino acids. Gapped positions were not included in the analyses. Bootstrap analyses were performed with 250 replicates, and all the resulting phylogenies were inspected to identify the most basal angiosperm lineage in cases where one clade was identified as sister to all other angiosperms with at least 50% bootstrap support.

Whereas alignment in many gapped regions was problematic, there were many regions where the homology of insertions or deletions could be assigned unambiguously. A separate data matrix of insertions and deletions was constructed from these regions, and parsimony analysis was performed on the binary data matrix.

#### Parametric Bootstrap Analyses

While much has been made of *Amborella* as the most basal clade in the angiosperm phylogeny (Mathews and Donoghue 1999; Parkinson, Adams, and Palmer 1999; Qiu et al. 1999; P. S. Soltis, D. E. Soltis, and Chase 1999; Barkman et al. 2000; Graham and Olmstead 2000;

Zanis et al. 2002; Borsch et al. 2003; Hilu et al. 2003), some analyses in these studies and others have found support for the placement of *Amborella* and Nymphaeales in a clade placed sister to all other angiosperms, whereas others have shown evidence for Nymphaeales alone as the most basal clade in the angiosperm phylogeny. Barkman et al. (2000) first showed that the placement of *Amborella* relative to Nymphaeales could vary among data sets and methods of phylogenetic reconstruction and predicted that incongruence among phylogenies estimated using different methods would increase as additional sequence data were added for each taxon in genome-scale analyses due to the effects of long-branch attraction (fig. 5 of Barkman et al. 2000). Following the comprehensive investigation of Zanis et al. (2002), we used parametric bootstrap analyses to explore whether long-branch attraction might be responsible for the observed conflict and determine if any of the four hypotheses illustrated in figure 1 could be rejected given the large number of chloroplast genome sequences in our study.

ML phylogenies were estimated in PAUP\* with constraint trees corresponding to one of four hypothesized lineages as sister to all other extant angiosperms: (1) *Amborella*, (2) an *Amborella* + Nymphaeales clade, (3) Nymphaeales, and (4) monocots (fig. 1). Parameter values for the HKY model with invariant sites and among-site rate variation were estimated in PHYML as described above. These parameter values and each of the four ML phylogenies shown in figure 1 were used to generate 200 simulated data sets with Seq-Gen version 1.3 (Rambaut and Grassly 1997). MP and NJ analyses were performed on all simulated data sets in PAUP\*, and ML analyses were performed using PHYML with five starting trees as described above. Model parameters for the ML and NJ analyses matched those used to simulate the data. The frequencies with which hypothesis a, b, c, or d was observed in the estimated phylogenies were calculated for each set of simulations and each phylogenetic methodology.

#### Dating Nodes

If some speciation events that gave rise to basal angiosperm lineages were separated by just a few million years, it may be difficult to resolve these events even with the large amount of data included in genome-scale studies with adequate taxon sampling. In order to explore timing of major events in angiosperm history, the ML phylogenies generated for each of the 200 nonparametric replicates were saved with branch lengths estimated to eight significant digits, and minimum ages were estimated for nodes on each phylogeny using the penalized likelihood method implemented in r8s (Sanderson 2003). The origin of the eudicots 125 MYA as evidenced by the appearance of tricolpate pollen in the fossil record (Crane, Friis, and Pederson 1995; Sanderson et al. 2004) was used as the fixed calibration point. In addition, minimum and maximum ages for the origin of the euphyllophytes were set at 380 and 410 MYA, respectively (Schneider et al. 2004), and minimum ages for the Poaceae and the most recent common ancestor (MRCA) of *Ginkgo* and *Pinus* were 55 MYA (Kellogg 2000) and 310 MYA (Schneider et al. 2004), respectively. Averages, medians, and standard errors were estimated for the

MRCA defined by the nodes in the ML tree across all 200 bootstrap trees.

## Results

### Monocots Are not Sister to All Other Extant Angiosperms

As Felsenstein (1978) famously deduced, there is a tendency for long branches to artifactually attract (referred to as “long-branch attraction”) in some phylogenetic analyses, and under these conditions, the probability of inferring the wrong phylogeny will increase with additional data due to statistical inconsistency. Hendy and Penny (1989) showed that even when rates of evolution are constant among lineages, long-branch attraction may confound parsimony analyses with more than four taxa and trees with variable terminal branch lengths. Because the out-group is almost always a long branch, this often manifests as the longer branch in-group taxa being drawn to the base of the tree even when this is not the correct relationship (Philippe and Laurent 1998). In such instances, adding taxa to interrupt the longest branches can help parsimony to converge on the correct phylogeny (Hendy and Penny 1989).

By adding nongrass monocots and another gymnosperm to the analysis, we interrupted two of the longest branches (fig. 3). As a result, all MP and ML analyses placed branching points for *Amborella* and Nymphaeales at the base of the angiosperm phylogeny with strong support (fig. 4). The most parsimonious nucleotide-based phylogeny constrained to place monocots sister to all other angiosperms was 242 steps longer than the unconstrained MP phylogeny (fig. 4B). A Wilcoxon signed ranks test showed this difference to be highly significant ( $P < 0.0001$ ). ML analyses with among-site rate variation placed *Amborella* + Nymphaeales sister to all other angiosperms (fig. 4A), whereas the ML analysis without rate variation placed *Amborella* as sister to all extant angiosperms (supplementary fig. 1A and B, see *Supplementary Material*). The log-likelihood score for the optimal HKY- $\Gamma$  + I likelihood tree in an analysis constrained to place monocots sister to all other angiosperms ( $-\ln(L) = 318,423.61$ ) was 310.73 points greater than the score for the unconstrained ML tree ( $-\ln(L) = 318,112.88$ ; figs. 1B and 4A). Based on the Shimodaira and Hasegawa (S-H; 1999) test, this difference in likelihood scores was highly significant ( $P < 0.0001$ ). The NJ analyses with among-site rate variation found strong support for an *Amborella* + Nymphaeales clade sister to all other angiosperms (fig. 4C), but the NJ analysis under the simpler model without variation in rates among sites placed the eudicots as sister to the remaining angiosperms (supplementary fig. 1C, see *Supplementary Material*). NJ analyses with the simple HKY model performed after removing the most distant out-group taxa, *Physcomitrella*, *Marchantia*, and *Psilotum*, placed an *Amborella* + water lilies clade as sister to the remaining angiosperms (supplementary fig. 1D, see *Supplementary Material*). ML and NJ analyses performed on the nucleotide data matrix using the more complicated GTR- $\Gamma$  + I model gave the same topologies shown in figure 4 with bootstrap values  $\pm 2\%$  relative to the results of the HKY- $\Gamma$  + I analysis (data not shown).




FIG. 3.—Comparison of unrooted MP phylogenies estimated from amino acid alignments with the taxon set analyzed by Goremykin et al. (2004, A) and in this study (B) reveals the long branch leading to grasses.

Analyses of the amino acids and first and second codon positions also placed *Amborella* and Nymphaeales at the base of the angiosperm phylogeny with strong support in the MP and ML analyses (supplementary figs. 2 and 3,

respectively, see *Supplementary Material*). NJ analyses of the first and second positions gave the same topology as the third codon position analysis, but the amino acid analysis placed eudicots as sister to the remaining angiosperms giving




FIG. 4.—Single MP (A), ML (B), and NJ (C) phylogenies estimated from the 61-gene nucleotide alignment are shown. Most nodes on each phylogeny were recovered in 100% of the bootstrap replicates, and only values <100% are shown for each node. All analyses place *Amborella* and the water lilies as basal lineages in the angiosperm phylogeny.

weak support for a (monocots (*Calycanthus*, (*Amborella*, Nymphaeales))) clade (supplementary figs. 2 and 3, respectively, see *Supplementary Material*). After removing *Physcomitrella*, *Marchantia*, and *Psilotum* from the amino acid alignment, NJ analyses recovered an *Amborella* + water lilies clade as sister to all other angiosperms. The placement of *Calycanthus*, which was poorly supported in all but two NJ analyses, differed among all three phylogenetic methods in nucleotide analyses (fig. 4). Whereas bootstrap support for the placement of *Calycanthus* was  $\leq 65\%$  in all MP and ML analyses, *Calycanthus* was placed sister to the monocots with support values of 100% and 90% in NJ analyses of the ungapped nucleotide and first and second codon position alignments, respectively. Despite high support values in these NJ analyses, additional sequences sampled from magnoliid orders other than Laurales (Magnoliales, Piperales, and Canellales), a member of the Chloranthaceae, and *Ceratophyllum* will be necessary to have any hope of resolving the relationships among these taxa, the monocots and eudicots, with confidence.

NJ analyses of LogDet distance matrices estimated from nucleotide and amino acid alignments assuming no among-site rate variation, six rate classes, and invariant sites (ML estimate of 26%) plus six additional rate classes all produced the same topology placing the core eudicots sister to a clade with all other eudicots including the eudicot *Ranunculus* (supplementary fig. 4, see *Supplementary Material*). The monophyly of the eudicots including the ranunculids as sister to the core eudicots is well established (Judd and Olmstead 2004), so this topology is quite unlikely to represent the true phylogeny. Moreover, analyses of nucleotide alignments including only seed plants gave strong support for an *Amborella* + water lilies clade as sister to the remaining angiosperms (supplementary fig. 4E and F, see *Supplementary Material*). The amino acid analysis of the seed plants, however, placed eudicots (including *Ranunculus*) sister to the remaining angiosperms (supplementary fig. 4G and H, see *Supplementary Material*). These results are noteworthy as they add to a growing set of examples underscoring the need for deeper understanding of how covarion/covariotide evolution and other forms of heterotachy can be diagnosed and modeled in phylogenetic analyses (e.g., Lockhart et al. 1998, 1999; Huelsenbeck 2002; Lopez, Casane, and Philippe 2002; Kolaczkowski and Thornton 2004; Phillips, Delsuc, and Penny 2004; Ané et al. 2005; Martin et al. 2005; Spencer, Susko, and Roger 2005).

In general, the relationships common to all trees in figure 4 were found in all ML trees and most NJ trees inferred from nucleotide and amino acid alignments given a variety of substitution models. Most of the exceptions seen in the NJ trees were only found in analyses that included distantly related out-group taxa. Most importantly, conflicts among topologies derived using different phylogenetic methods, substitution models, or taxon sets were usually supported by bootstrap values greater than 90%. This observation underscores how statistical inconsistency is a serious problem for phylogenomics that can only be diagnosed through comprehensive analyses using a variety of phylogenetic methods, substitution models, distance corrections, and taxon sets (see also Phillips, Delsuc, and Penny 2004).

#### *Amborella* or *Amborella* Plus Nymphaeales?

As has been described in previous studies (Barkman et al. 2000; Zanis et al. 2002; Stefanović, Rice, and Palmer 2004), the relationship between *Amborella* and the Nymphaeales relative to the rest of the angiosperms depended on the method of analysis. In our MP analyses, *Amborella* was placed sister to all other extant angiosperms with high support (fig. 4B). A tree constrained to have a Nymphaeales + *Amborella* clade was 83 steps longer than the MP tree, and this difference was found to be significant in a Wilcoxon signed ranks test ( $P = 0.0007$ ). Examination of the characters potentially supporting an *Amborella* alone versus a Nymphaeales + *Amborella* clade showed that characters were distributed evenly across the 61-gene alignment. However, despite strong support for the basal branching point for *Amborella* under MP, the NJ and ML analyses unite Nymphaeales with *Amborella* in a clade that is sister to the rest of the angiosperms. The ML analysis gave moderate support for this relationship (fig. 4B), whereas the NJ analysis provided strong support (fig. 3C). The likelihood score ( $-\ln(L) = 318,117.04$ ) for a tree constrained to place *Amborella* as sister to all other angiosperms was not significantly different than that of the ML tree ( $P = 0.601$ ; scores shown in fig. 1). Furthermore, ML analyses of the amino acid alignments provided weak support for *Amborella* as sister to the remaining angiosperms (supplementary fig. 3, see *Supplementary Material*). The tree placing the water lilies as sister to the rest of the angiosperms gave a significantly worse likelihood score than the ML phylogeny ( $P = 0.028$ ).

Most single-gene analyses did not resolve basal relationships among angiosperm lineages (table 2). In all character sets, however, when one angiosperm lineage was placed sister to the others, it was most often *Amborella* or *Amborella* + Nymphaeales (table 2). The monocots were not resolved as sister to the other angiosperms in any of the single-gene phylogenies, and Poaceae or Poales (Poaceae + *Typha*) were estimated as the most basal clade for only four genes in analyses of codon positions 1 and 2. A few of the single-gene analyses supported relationships that are clearly untenable (table 2). For example, a weakly supported Poaceae + *Oenothera* clade was placed sister to the rest of the angiosperms in the analysis of *clpP* nucleotides. The most basal angiosperm lineage inferred for some genes varied across the three character sets. For example, the *psbB* phylogenies estimated from all nucleotides, first and second codon positions and amino acids were core eudicots, Poales, and *Typha*, respectively.

The MP analysis of insertions and deletions also provided strong support for *Amborella* and Nymphaeales as branching from the most basal nodes in the angiosperm phylogeny (fig. 5 and supplementary tables 1 and 2, see *Supplementary Material*). A total of 116 potentially phylogenetically informative indels were included in the analysis. Four unambiguous indels, consisting of three insertions and one deletion mutation, supported the monophyly of all sampled angiosperms except *Amborella* and Nymphaeales (fig. 6). No indel characters were potentially supportive of grasses or monocots in the basal-most position. Twelve equally parsimonious trees (141 steps) were recovered in

**Table 2**  
**The Lineage(s) That Is Inferred to Be Most Basal in Angiosperms Was Ambiguous in Most Single-Gene Analyses**

Taxon <sup>1</sup>	Codon Positions 1 + 2 + 3	Codon Positions 1 + 2	Amino Acids
Poaceae + <i>Oenothera</i>	1 (clpP [80])		
Poaceae		3 (rpoB [50], psbK [54], rps3 [53])	2 (rpoC1 [86], rps12 [64])
<i>Typha</i>			1 (psbB [53])
Poales ( <i>Typha</i> + Poaceae)		1 (psbB [53])	
Eudicots	1 (rps2 [82])		
Core Eudicots	2 (psbB [53], psbD [54])		
<i>Oenothera</i>		1 (rps2 [50])	1 (rps2 [69])
<i>Spinacia</i>		1 (petD [53])	1 (rps15 [66])
<i>Amborella</i>	5 (atpE [61], atpF [56], psaA [59], rbcL [59], rpoA [86])	3 (atpE [58], rpoA [64], cemA [68])	2 (atpE [57], cemA [64])
Nymphaeales	1 (rpoB [69])		
Nymphaeales + <i>Amborella</i>	3 (rpoC1 [66], rpoC2 [74], rps4 [65])	2 (rpoC2 [73], matK [51])	1 (ccsA [53])
Unresolved	48	50	54
Total	61	61	61

NOTE.—The identities of basal lineages inferred with at least 50% bootstrap support in the MP analyses are shown for the nucleotide and amino acid alignments with all gapped positions removed. Bootstrap support (percent) for basal position indicated lineage is shown in square brackets for each gene.

the MP analysis of the indel characters. Half of the MP trees included an *Amborella* + Nymphaeales clade, and half placed *Amborella* alone as sister to all other angiosperm lineages. Each of these two hypotheses was supported by a single synapomorphy (fig. 6). The resolved portions of the indel phylogeny were identical to corresponding relationships inferred through comparisons of nucleotide and amino acid sequences.

A large number of the microstructural mutations distinguish grasses from all other plants in the study (fig. 5B), whereas other lineages have many fewer insertion-deletion mutations. This suggests that lineage-specific common

processes may have given rise to the enhanced rates of nucleotide substitution and indel evolution in the lineage leading to the Poaceae, and perhaps generally throughout the major lineages of land plant evolution.

#### Parametric Bootstrap Analysis

The parametric bootstrap analysis demonstrated that the results of the MP analysis may have been affected by long-branch attraction (Felsenstein 1978; Henny and Penny 1989). Whereas the ML and NJ analyses recovered the simulated topology in the vast majority of cases, the MP




FIG. 5.—Parsimony analysis of microstructural mutations in 61 coding regions supports *Amborella* and water lilies as the most basal lineages of angiosperms: (A) bootstrap analysis of 116 parsimony-informative insertion and deletion characters, (B) phylogram of 1 of the 12 most parsimonious trees (141 steps) with branch lengths drawn proportional to the number of inferred insertion and deletion mutations on each branch.




FIG. 6.—Six phylogenetically informative insertion-deletion mutations bearing on the position of *Amborella* and water lilies relative to other angiosperms. Each minialignment is identified by the gene name and position in the *Amborella* sequence of the first amino acid in the indel of interest. Four indels (A, B, C, and D) support the basal position of *Amborella* and water lilies; one indel (E) supports *Amborella* as the sole basal-most angiosperm lineage; one indel supports the monophyly of *Amborella* plus water lilies. Thus, characters E and F are consistent with A–D but conflict with regard to the relationship between *Amborella* and water lilies. No phylogenetically informative indel characters were observed that would have supported a basal position for grasses or monocots in the angiosperms.

analyses misidentified *Amborella* as sister to all other angiosperms in 21% of the cases where data were simulated under the *Amborella* + Nymphaeales phylogeny (table 3). When constrained to place monocots sister to the rest of the angiosperms, the ML phylogeny included a very short internal branch leading to the node where an *Amborella* + Nymphaeales clade diverged from the remaining dicot lineages (fig. 1D). Data sets generated under this hypothesis were especially problematic for MP and NJ analyses. Only ML recovered the correct tree in more than half of the analyses (table 3). In exploratory parametric bootstrap analyses where Nymphaeales was only represented by the *Nuphar* sequence, both NJ and MP analyses failed to recover the correct topology for more than 97% of the data sets simu-

lated under the *Amborella* + Nymphaeales phylogeny. The effect of adding *Nymphaea* to the analysis illustrates the strong influence of taxon sampling on phylogenetic reconstruction.

Molecular Clock Estimates

Molecular clock estimates for most angiosperm nodes in the ML topology were in line with recently published divergence dates estimated using a variety of procedures (Chaw et al. 2004; Davies et al. 2004; others reviewed in Sanderson et al. 2004; table 4; fig. 7). The median age estimates for the angiosperm crown group and the MRCA of all monocots, magnoliids, and eudicots were

**Table 3**  
**Parametric Bootstrap Results Show that Parsimony Analyses Are More Subject to Long-Branch Attraction than the Model-Based Likelihood and Neighbor-Joining Analyses When Data Sets are Simulated on Phylogenies B (*Amborella* + Nymphaeales basal clade) and C (Nymphaeales basal-most clade)**

Simulated Phylogeny <sup>a</sup>	Likelihood				Parsimony				Neighbor Joining			
	Amb	Amb + Nym	Nym	Monocot	Amb	Amb + Nym	Nym	Monocot	Amb	Amb + Nym	Nym	Monocot
A	<b>100</b>	—	—	—	<b>100</b>	—	—	—	<b>100</b>	—	—	—
B	—	<b>100</b>	—	—	21	<b>79</b>	—	—	—	<b>100</b>	—	—
C	—	—	<b>100</b>	—	10	—	<b>90</b>	—	1	1	<b>98</b>	—
D <sup>b</sup>	—	1	—	<b>98</b>	—	48	—	<b>4</b>	—	29	—	<b>41</b>

NOTE.—Parsimony and neighbor joining performed poorly when data were simulated on the ML tree forcing monocots as sister to all other angiosperms. Rate of recovering the simulated topology is shown in bold for each reconstruction method and simulated phylogeny. Amb, *Amborella*; Nym, Nymphaeales.

<sup>a</sup> Input phylogenies for the simulations are shown in figure 1.

<sup>b</sup> Rows do not sum to one for tree D because other angiosperm rootings were observed in the bootstrap trees, including monocots + *Amborella* + Nymphaeales, *Calycanthus* + monocots + *Amborella* + Nymphaeales and *Calycanthus* as sister to remaining angiosperms.

**Table 4**  
**Estimated Ages (in millions of years) for Nodes Labeled**  
**in Figure 6**

Node Label	Description	Constraint	Estimated Age
A	Euphyllophytes	380 minimum 401 maximum	410
b	Seed plants		334
c	<i>Pinus</i> + <i>Ginkgo</i>	310 minimum	310
d	Angiosperms	132 minimum	161 (158–165)
e	Nymphaeales/ <i>Amborella</i>		136 (134–165)
f	Nymphaeaceae s.s.		22 (21–23)
g	Magnoliids/monocots/eudicots		145 (143–147)
h	Monocots		133 (131–135)
i	Asparagales/commelinids		117 (116–118)
j	Poales		107 (106–109)
k	Core Poaceae	55 minimum	55
L	Eudicots	125 fixed	125
m	Core eudicots		113
n	Rosids		108 (108–109)
o	Faboideae		61 (61–62)
p	Asterids/Caryophyllales		105
q	Solanaceae		12

NOTE.—The median estimates are shown with the range of observed estimates for 200 bootstrap replicates. All estimates were within 1 Myr for the nodes with no range shown.

161 MYA and 145 MYA, respectively. The bootstrap distribution of divergence time estimates was unimodal with low standard error estimates for most nodes with a notable exception (table 4). Age estimates for the MRCA of *Amborella* and the Nymphaeales had two modes, corresponding to bootstrap replicate ML topologies with and without an *Amborella* + Nymphaeales clade. Trees with *Amborella* and the water lilies forming a clade gave a median age estimate of 135 MYA for their MRCA, whereas the median age estimate was 161 MYA for the 68 bootstrap replicates where *Amborella* was found to be sister to all other angiosperms.

The range of age estimates calculated from 200 bootstrap replicates is shown in table 4. It should be noted that whereas the small errors in our branch-lengths estimates follow legitimately from the large number of nucleotide positions included in our analyses, the errors on our divergence date estimates are artificially low given that our dating analysis was done under the assumption that calibration points for the MRCAs of all eudicots is known without error (see Graur and Martin 2004). The appearance of tricolpate pollen in the fossil record at the Barremian-Aptian boundary 125 MYA provides a minimum age for the MRCA of all extant eudicots.

## Discussion

Phylogenetic analyses of plant plastid genomes are providing new insights into the evolution of gene order (Cosner, Raubeson, and Jansen 2004), lineage-specific substitution rates and patterns (Ané et al. 2005), and factors influencing genome-scale phylogenetic inference (Lockhart et al. 1999; Goremykin et al. 2003, 2004; Soltis et al. 2004b; Stefanović, Rice, and Palmer 2004; Lockhart and Penny 2005; Martin et al. 2005). Alignment and orthology assignments are straightforward for the majority of coding regions, making plastid genomes ideal for phylogenetic reconstruction and studies of molecular evolution. Whole




FIG. 7.—The ML phylogeny inferred from the nucleotide analysis with labeled nodes corresponding to the minimum divergence time estimates shown in table 4.

genome sequencing of plastid genomes provide copious data for testing hypothesized organismal relationships, comparing models of molecular evolution, and developing analytical methodologies. At this point, however, with few plastid genomes available for analysis, care must be taken to avoid being misled by the results of some analyses. With nearly 40,000 sites in our ungapped nucleotide alignments of 61 genes, any method that is susceptible to statistical inconsistency may be affected by long-branch attraction.

Our results lead us to reject all but two hypotheses concerning the basal-most extant angiosperm lineages. The ML analyses provide weak support for *Amborella* and Nymphaeales as a clade sister to all other angiosperms, but the more popular hypothesis placing *Amborella* alone as sister to all other angiosperms could not be rejected. The monocots clearly constitute a slightly younger clade in the angiosperm phylogeny. We estimate that the divergence of monocot and eudicot lineages occurred only 16 Myr after the MRCA of extant angiosperms (table 4). To put this result into context, the estimated dates for the youngest nodes on the ML phylogeny, *Nicotiana/Atropa* ( $10.56 \pm 0.03$ ) and *Zea/Saccharum* ( $8.87 \pm 0.01$ ), are just over half this age difference.

Despite the short internodes at the base of the angiosperm phylogeny, branching order can be resolved correctly with sufficient data. We conclude, however, that a lineage-specific increase in nucleotide substitution rates on the branch leading to the grasses and incomplete taxon

sampling in the monocots confounded the analyses of Goremykin et al. (2003, 2004), resulting in the inference of the grasses sister to all other angiosperms in all analyses that did not include among-site rate variation. The earlier studies did place the branching point for *Amborella* or *Amborella* + Nymphaeales at the basal angiosperm node when ML analyses included a correction of among-site rate variation (Goremykin et al. 2003, 2004; Stefanović, Rice, and Palmer 2004). With additional monocots included in the data matrix (table 1), we find that all MP and ML analyses of our 61-gene alignments (both nucleotides and amino acids) placed the branching point(s) for *Amborella* and the Nymphaeales at the base of the angiosperm phylogeny with strong support. Some of the distance-based NJ analyses placed core eudicots or eudicots as sister to the remaining angiosperms. However, all but the LogDet analyses of the amino acid alignment provided strong support for an *Amborella* + water lilies clade as sister to all other extant angiosperms when the analyses were restricted to seed plants (supplementary figures 1 and 4, see *Supplementary Material*). The basis of differences in the results of LogDet analyses performed on the nucleotide and amino acid alignments deserves further investigation.

As has been described previously (e.g., Eyre-Walker and Gaut 1997), the substitution rate for chloroplast genes is accelerated both within the grasses and on the branch leading to the most common ancestor of maize, rice, and wheat (fig. 3). The earlier work found an increase in synonymous substitution after the divergence of the grasses and the palms (Areaceae, Arecales). The phylograms shown in figures 1, 3, and 4 suggest that the rate acceleration occurred within the Poales, after divergence from the MRCA of *Typha* (Typhaceae, Poales) and the grasses.

The phylogenetic position of the Nymphaeales relative to *Amborella* and the remaining angiosperms remains unresolved. While the parsimony analysis suggests strong support for *Amborella* as sister to all other angiosperms, parametric bootstrap analyses performed here and in a previous study (Zanis et al. 2002) lead us to interpret the parsimony results cautiously. At the same time, the strong support for a *Amborella* + Nymphaeales clade observed in the NJ analysis is tempered by the moderate support for this clade in the ML analysis and the miniscule, nonsignificant difference in likelihood scores between the topologies *A* and *B* in figure 1. Whether the topology represented in figure 1(*A* or *B*) is correct, the molecular clock analysis suggests that the Nymphaeales lineage diverged from a sister lineage leading either to *Amborella* or to all other angiosperms some 25 Myr after the MRCA of all extant angiosperms.

Previous studies have reached different conclusions concerning the relationship of *Amborella* and the Nymphaeales at the base of the angiosperm phylogeny. The initial identification of *Amborella*, the Nymphaeales, and the Austrobaileyales as the most basal lineages of extant angiosperms (Mathews and Donoghue 1999; Parkinson, Adams, and Palmer 1999; Qiu et al. 1999; P. S. Soltis, D. E. Soltis, and Chase 1999; Graham and Olmstead 2000) was a landmark event in molecular systematics. Although *Amborella* was favored as sister to all other extant angiosperm lineages in these seminal multigene studies (hypothesis a, fig. 1A), a hypothesis placing *Amborella* with Nymphaeales in a clade

sister to the remaining angiosperm lineages (hypothesis b, fig. 1B) could not be rejected (Parkinson, Adams, and Palmer 1999; Mathews and Donoghue 2000; Qiu et al. 2000). Barkman et al. (2000) favored the *Amborella* + Nymphaeales basal clade hypothesis after performing a series of MP, ML, and NJ analyses on partitioned and complete multigene data sets, both before and after applying a controversial “noise-reduction” screen designed to identify and remove sites that may obscure phylogenetic signal (Lyons-Weiler, Hoelzer, and Tausch 1996). However, some of their analyses provided strong support for hypothesis a, and many gave weak to moderate support for either hypothesis. They used a nonparametric bootstrap resampling procedure to test their prediction that as larger amounts of data were gathered from genes with similar evolutionary dynamics to those sampled the support for the method-dependent conflict would grow increasingly strong due to statistical inconsistency.

Zanis et al. (2002) also found that inference of the relationship between *Amborella* and Nymphaeales was dependent on data partition and phylogenetic method. In general agreement with the previous studies of DNA sequence data, MP and nuclear ribosomal genes offered the strongest support for hypothesis a, while ML analyses of protein-coding genes and genes sampled from the chloroplast and mitochondrial genomes gave weak support for hypothesis b. MP and ML analyses of the combined data set gave rather strong support for hypothesis a, but hypothesis b could not be rejected in a likelihood ratio test. This result is very similar to our finding of 63% bootstrap support for hypothesis b while hypothesis a cannot be rejected in the S-H test. Zanis et al. found as we did that parametric bootstrap analyses demonstrated bias in MP reconstruction toward recovery of *Amborella* as sister to all other extant angiosperms.

Despite the very large alignments analyzed for this study, we are still unable to conclusively reject either hypothesis a or b (fig. 1). Accurate phylogenetic resolution can generally be achieved more efficiently when taxa can be added to break long branches on the phylogeny (Graybeal 1998; Pollock et al. 2002; Zwickl and Hillis 2002; Hillis et al. 2003). This generalization, however, may not always apply to the resolution of branching order among basal lineages (Simmons and Miya 2004). Recent studies with extensive taxon sampling (P. S. Soltis, D. E. Soltis, and Chase 1999; Zanis et al. 2002 [data set 2]; Hilu et al. 2003) have supported hypothesis a, with moderate to high levels of support in MP analyses. It is not clear, however, whether increased taxon sampling was sufficient in these studies to interrupt the long branches responsible for the bias observed in the parametric bootstrap analyses performed here and by Zanis et al. (2002).

The results presented by Goremykin et al. (2003, 2004) demonstrate how incomplete taxon sampling can result in strong support for conflicting topological relationships. Graham and Olmstead (2000) had previously shown how sampling among basal angiosperm lineages can influence phylogenetic reconstruction of branching order. Their MP analysis found strong support (96%) for Nymphaeales as sister to all other angiosperms (e.g., fig. 1C) when the order was represented by *Cabomba*

(Cabombaceae) alone, but when *Nymphaea* and *Cabomba* were included in the analysis, *Amborella* moved to the base of the angiosperm phylogeny with moderate support (69%). When we added data from our six new plastid genome sequences one at a time to the 61-gene nucleotide matrices, we found that the addition of monocots *Typha* and *Yucca* changed the strongly supported position of the grasses in MP phylogenies (supplementary fig. 5, see *Supplementary Material*). As was found by Stefanović, Rice, and Palmer (2004), the grasses and *Acorus* were placed as successive sister lineages to the rest of the angiosperms in MP analyses when *Acorus* and the grasses were the only monocots included in the analysis (supplementary fig. 5F, see *Supplementary Material*). As reported by Goremykin et al. (2003, 2004), we found that ML analyses including variation across sites placed lineages leading to *Amborella* and the water lilies at the base of the angiosperm phylogeny irrespective of taxon set (supplementary fig. 6, see *Supplementary Material*).

Due to extinctions, taxa are not available to reduce the length of the critical branch separating the angiosperms and gymnosperms or the terminal branch leading to *Amborella*. It is possible that the combination of these long branches and the short internode subtending the branching point for the Nymphaeales and its sister lineage (*Amborella* [hypothesis b] or the rest of the extant angiosperms [hypothesis a]) may not allow us to conclusively reject hypothesis a or b. The possibility of long-branch attraction under these circumstances is expected to be especially problematic in analyses using only rapidly evolving coding (Hilu et al. 2003) and noncoding (Borsch et al. 2003) sequences. The addition of species within the Cabombaceae (Nymphaeales), the Austrobaileyales, and magnoliids to the 61-gene data set, however, may lead to resolution of the relationships of *Amborella*, the Nymphaeales, and the remaining extant angiosperms. Aside from the improvement in phylogenetic analyses based on nucleotide and amino acid substitutions, the addition of these and out-group taxa to the 61-gene data matrix will likely improve alignment of gapped regions and increase the number of unambiguously scored indel characters.

Although all the prior phylogenetic analyses of whole plastid genome sequence have focused on DNA or protein sequence analyses, Graham and Olmstead (2000) and Graham et al. (2000) showed that solid phylogenetic inference can be derived from careful characterization of insertion and deletion mutations in chloroplast genomes. Although much less numerous than base substitutions within most coding regions, our results agree with those of Graham et al. (2000), supporting the evidence that these characters have much lower homoplasy than base substitutions and may provide a special collection of evidence bearing on branching events that are otherwise challenging to resolve due to phylogenetic artifacts such as long-branch attraction (Graham et al. 2000; Rokas and Holland 2000). The unambiguous placement of *Amborella* and Nymphaeales as the most basal angiosperm lineages was resolved with our indel matrix even with the limited taxon sampling employed by Goremykin et al. (2003, 2004) (supplementary fig. 7, see *Supplementary Material*). The fact that the microstructural data does not appear to be affected by the same long-branch

attraction problems is noteworthy because it is clear (fig. 5B) that the rate of indel evolution is also dramatically increased in the lineage leading to the Poaceae. This result should motivate an extensive effort to identify more microstructural characters in these genomes.

We noted in our alignments many other regions that were rich in indel mutations but where unambiguous character assignment was not yet possible in our judgment. It is likely that as additional genomes are sequenced alignment of these difficult regions will improve, allowing the coding of many additional microstructural characters. Many of the lineage-specific indels that were ignored in this study should then emerge as synapomorphies among additional taxa.

These additions should also help resolve the other difficult nodes in the phylogenies involving the branching order of the magnoliids, monocots, and eudicots. Relationships among these three clades, *Ceratophyllum*, and Chloranthaceae have not been well resolved in previous studies, but the high support for the monophyly of the magnoliids (Magnoliales, Laurales, Canellales, and Piperales) in the 17-gene analysis of Graham and Olmstead (2000) suggests that whole chloroplast genome sequences could provide enough phylogenetically informative nucleotide variation to clarify relationships among these taxa.

The eudicots (tricolpates) comprise roughly 64% of angiosperm species diversity (Judd and Olmstead 2004). While many nodes within the phylogeny for the group are well supported, rapid diversification has made resolution of some nodes quite difficult. Resolution of the relationships among the major core eudicot lineages, including the Caryophyllales, rosids and asterids, has been particularly recalcitrant. The moderate to high bootstrap support observed for a clade joining the spinach lineage (Amaranthaceae, Caryophyllales) with *Atropa* and *Nicotiana* (Solanaceae, Solanales, euasterid I) should be interpreted cautiously. The chloroplast genome of *Panax* (Araliaceae, Apiales, euasterid II) has recently been published (Kim and Lee 2004), and its inclusion in the 61-gene data set results in slightly reduced bootstrap support for a Caryophyllales + asterid clade (supplementary fig. 8, see *Supplementary Material*).

Over the last decade, advances in our understanding of phylogenetic relationships among extant angiosperms (Chase 2004; Judd and Olmstead 2004; P. S. Soltis and D. E. Soltis 2004) have provided an improved framework for comparative analyses designed to elucidate the evolution of important features ranging from endosperm development (Williams and Friedman 2002) to MADS box gene evolution (Becker and Theissen 2003; Litt and Irish 2003; Kim et al. 2004; Kramer, Jaramillo, and Di Stillo 2004; Stellari, Jaramillo, and Kramer 2004) to the evolution of floral perianth organization (Zanis et al. 2003; Soltis et al. 2004a). Just as favored evolutionary scenarios had to be abandoned with the demise of the anthophyte hypothesis (Goremykin et al. 1996), inferences drawn in these and many other comparative studies would have had to be re-examined if the position of the monocots recovered in the phylogenies of Goremykin et al. (2003, 2004) were supported in subsequent studies. This study and others (D. E. Soltis and P. S. Soltis 2004; Stefanović, Rice, and

Palmer 2004), however, have tested and rejected the hypothesized position of monocots sister to all other angiosperm lineages (Goremykin et al. 2003, 2004).

As genome-scale sequencing and phylogenetic analyses become more common, the possible influence of long-branch attraction must be seriously considered in any interpretation of the resulting phylogenies. Phylogenies based on many genes sampled from a few model species will be especially susceptible to long-branch attraction (Soltis et al. 2004b; Philippe, Lartillot, and Brinkmann 2005). We contend that genome-scale phylogenetic studies can avoid misinterpretation of artifactual results by employing parametric bootstrap analyses (e.g., Sanderson et al. 2000; Zanis et al. 2002), multiple reconstruction methods (MP, ML, NJ, Bayesian), a variety of models of molecular evolution (e.g., Phillips, Delsuc, and Penny 2004; Stefanović, Rice, and Palmer 2004), consideration of variation in substitution patterns among lineages (Lockhart et al. 1998, 1999; Huelsenbeck 2002; Lopez, Casane, and Philippe 2002; Kolaczowski and Thornton 2004; Phillips, Delsuc, and Penny 2004; Ané et al. 2005; Martin et al. 2005; Spencer, Susko, and Roger 2005), taxon subsampling (Graham and Olmstead 2000; D. E. Soltis and P. S. Soltis 2004), and analyses of multiple data partitions (e.g., Barkman et al. 2000; Zanis et al. 2002). Inconsistencies among results derived from different approaches, models, or data sets should be examined and explained rather than ignored.

## Supplementary Material

Supplementary figures are available at *Molecular Biology and Evolution* online (<http://www.mbe.oxfordjournals.org/>)

SUPPLEMENTARY FIG. 1.—ML and NJ phylogenies recovered using the HKY substitution model without correction for among-site variation. Comparison of ML (A and B) and NJ phylogenies (C and D) estimated from alignments including euphyllophytes (A and C) or just seed plant sequences (B and D) shows that the NJ analyses run under the simple HKY model are influenced by the inclusion or exclusion of distant out-group sequences.

SUPPLEMENTARY FIG. 2.—ML (A), MP (B), and NJ (C) phylogenies with bootstrap values from analyses of first and second codon positions in the nucleotide alignments are very similar to those estimated with all three codon positions (fig. 4). The bootstrap value increased for the water lilies + *Amborella* clade in the ML analysis, and the poorly supported placement of *Calycanthus* relative to the eudicots and monocots changed in the MP analysis. ML and NJ analyses were performed using the HKY +  $\Gamma$  + I model as described in text. Bootstrap values are not shown for branches with 100% support.

SUPPLEMENTARY FIG. 3.—The results of ML (A) and NJ (C) analyses of amino acid alignment differ slightly from those estimated with the complete nucleotide alignment (fig. 4). The ML analysis (JTT +  $\Gamma$  + I) returns poor support of *Amborella* as sister to all other angiosperms, and the NJ analysis (JTT +  $\Gamma$ ) places eudicots as sister to the remaining angiosperms. An NJ analysis restricted to the seed plants returns relationships identical to those for seed plants shown in figure 4(C).

SUPPLEMENTARY FIG. 4.—Phylogenies resulting from analyses performed on the complete nucleotide and amino acid alignments (ungapped) using LogDet corrected distances place as the core eudicots as sister all other angiosperms including *Ranunculus* (A–D). Whereas analyses restricted to the seed plant nucleotide alignment recover relationships identical to those for seed plants shown in figure 4(C) (E and F), the eudicots are placed sister to the remaining angiosperms in analyses of the amino acid alignment for seed plants (G and H). Topologies are identical for LogDet analyses performed with (B, D, F, and H) and without (A, C, E, and G) variation rates across sites (see text).

SUPPLEMENTARY FIG. 5.—Phylogenies from MP analyses adding *Ginkgo* (A), *Nuphar* (B), *Ranunculus* (C), *Acorus* (D), *Yucca* (E), and *Typha* (F) one at a time to a 61-gene nucleotide alignment of previously available plastid genomes.

SUPPLEMENTARY FIG. 6.—Phylogenies from ML analyses adding *Ginkgo* (A), *Nuphar* (B), *Ranunculus* (C), *Acorus* (D), *Yucca* (E), and *Typha* (F) one at a time to a 61 gene nucleotide alignment of previously available plastid genomes. The HKY +  $\Gamma$  + I substitution model was used in all analyses.

SUPPLEMENTARY FIG. 7.—Parsimony bootstrap consensus trees of indel characters, using taxon sets from (A) Goremykin et al. (2003) and (B) Goremykin (2004). For analysis A, one MP tree was obtained (115 steps; consistency index (CI) = 0.9111; RC = 0.8856), with *Amborella* as the first branching angiosperm, while B obtained two MP trees at 127 steps (CI = 9134; RC = 0.8560), one with *Amborella* and one with *Amborella* + *Nymphaea* as the earliest angiosperm branch.

SUPPLEMENTARY FIG. 8.—Bootstrap consensus phylogenies for ML, MP and NJ analyses of nucleotide alignment including 61 genes from *Panax schinseng* plastid genome sequence (Kim and Lee 2004) are consistent with those shown in figure 4. All analyses were performed as described in text.

SUPPLEMENTARY DATA MATRIX 1. Nexus file with nucleotide alignment. <http://chloroplast.cbio.psu.edu/>

SUPPLEMENTARY DATA MATRIX 2. Nexus file with amino acid alignment. <http://chloroplast.cbio.psu.edu/>

SUPPLEMENTARY DATA MATRIX 3. Nexus file with microstructural characters. <http://chloroplast.cbio.psu.edu/>

## Acknowledgments

We wish to thank Sean Graham, Jeff Palmer, Bill Martin, David Penny, Doug Soltis, and an anonymous reviewer for helpful discussions concerning this work. Funding was provided by the National Science Foundation grants DEB-0120709, DEB-0234417 and DBI-0115684.

## Literature Cited

- Ané, C., J. G. Burleigh, M. M. McMahon, and M. J. Sanderson. 2005. Covarion structure in plastid genome evolution: a new statistical test. *Mol. Biol. Evol.* **22**:914–924.
- Asano, T., T. Tsudzuki, S. Takahashi, H. Shimada, and K. Kadowaki. 2004. Complete nucleotide sequence of the sugarcane (*Saccharum officinarum*) chloroplast genome:

- a comparative analysis of four monocot chloroplast genomes. *DNA Res.* **11**:93–99.
- Barkman, T. J., G. Chenery, J. R. McNeal, J. Lyons-Weiler, W. J. Ellisens, G. Moore, A. D. Wolfe, and C. W. dePamphilis. 2000. Independent and combined analyses of sequences from all three genomic compartments converge on the root of flowering plant phylogeny. *Proc. Natl. Acad. Sci. USA* **97**:13166–13171.
- Becker, A., and G. Theissen. 2003. The major clades of MADS-box genes and their role in the development and evolution of flowering plants. *Mol. Phylogenet. Evol.* **29**:464–489.
- Borsch, T., K. W. Hilu, D. Quandt, V. Wilde, C. Neinhuis, and W. Barthlott. 2003. Noncoding plastid trnT-trnF sequences reveal a well resolved phylogeny of basal angiosperms. *J. Evol. Biol.* **16**:558–576.
- Chase, M. W. 2004. Monocot relationships: an overview. *Am. J. Bot.* **91**:1645–1655.
- Chaw, S. M., C. C. Chang, H. L. Chen, and W. H. Li. 2004. Dating the monocot-dicot divergence and the origin of core eudicots using whole chloroplast genomes. *J. Mol. Evol.* **58**:424–441.
- Cosner, M. E., L. A. Raubeson, and R. K. Jansen. 2004. Chloroplast DNA rearrangements in Campanulaceae: phylogenetic utility of highly rearranged genomes. *BMC Evol. Biol.* **4**:27.
- Crane, P. R., E. M. Friis, and K. R. Pederson. 1995. The origin and early diversification of angiosperms. *Nature* **374**:27–33.
- Darwin, C. 1903. Letter to J. D. Hooker. *In* F. Darwin and A. C. Seward, eds. *More letters of Charles Darwin*, Vol. 2. John Murray, London.
- Davies, T. J., T. G. Barraclough, M. W. Chase, P. S. Soltis, D. E. Soltis, and V. Savolainen. 2004. Darwin's abominable mystery: insights from a supertree of the angiosperms. *Proc. Natl. Acad. Sci. USA* **101**:1904–1909.
- Ewing, B., and P. Green. 1998. Base-calling of automated sequencer traces using phred. II. Error probabilities. *Genome Res.* **8**:186–194.
- Ewing, B., L. Hillier, M. C. Wendl, and P. Green. 1998. Base-calling of automated sequencer traces using phred. I. Accuracy assessment. *Genome Res.* **8**:175–185.
- Eyre-Walker, A., and B. S. Gaut. 1997. Correlated rates of synonymous site evolution across plant genomes. *Mol. Biol. Evol.* **14**:455–460.
- Felsenstein, J. 1978. Cases in which parsimony and compatibility methods will be positively misleading. *Syst. Zool.* **27**:401–410.
- . 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**:783–791.
- . 2004. PHYLIP (phylogeny inference package). Version 3.6. Distributed by the author, Department of Genome Sciences, University of Washington, Seattle.
- Gordon, D., C. Abajian, and P. Green. 1998. Consed: a graphical tool for sequence finishing. *Genome Res.* **8**:195–202.
- Goremykin, V., V. Bobrova, J. Pahnke, A. Troitsky, A. Antonov, and W. Martin. 1996. Noncoding sequences from the slowly evolving chloroplast inverted repeat in addition to *rbcL* data do not support gnetalean affinities of angiosperms. *Mol. Biol. Evol.* **13**:383–396.
- Goremykin, V. V., K. I. Hirsch-Ernst, S. Wolf, and F. H. Hellwig. 2003. Analysis of the *Amborella trichopoda* chloroplast genome sequence suggests that *Amborella* is not a basal angiosperm. *Mol. Biol. Evol.* **20**:1499–1505.
- . 2004. The chloroplast genome of *Nymphaea alba*: whole-genome analyses and the problem of identifying the most basal angiosperm. *Mol. Biol. Evol.* **21**:1445–1454.
- Graham, S. W., and R. G. Olmstead. 2000. Utility of 17 chloroplast genes for inferring the phylogeny of the basal angiosperms. *Am. J. Bot.* **87**:1712–1730.
- Graham, S. W., P. A. Reeves, A. C. E. Burns, and R. G. Olmstead. 2000. Microstructural changes in noncoding chloroplast DNA: interpretation, evolution, and utility of indels and inversions in basal angiosperm phylogenetic inference. *Int. J. Plant Sci.* **161**:S83–S96.
- Graur, D., and W. Martin. 2004. Reading the entrails of chickens: molecular timescales of evolution and the illusion of precision. *Trends Genet.* **20**:80–86.
- Graybeal, A. 1998. Is it better to add taxa or characters to a difficult phylogenetic problem? *Syst. Biol.* **47**:9–17.
- Guindon, S., and O. Gascuel. 2003. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst. Biol.* **52**:696–704.
- Hasegawa, M., H. Kishino, and T. Yano. 1985. Dating of the human-ape splitting by a molecular clock of mitochondrial DNA. *J. Mol. Evol.* **22**:160–174.
- Hendy, M. D., and D. Penny. 1989. A framework for the quantitative study of evolutionary trees. *Syst. Zool.* **38**:297–309.
- Hillis, D. M., D. D. Pollock, J. A. McGuire, and D. J. Zwickl. 2003. Is sparse taxon sampling a problem for phylogenetic inference? *Syst. Biol.* **52**:124–126.
- Hilu, K. W., T. Borsch, K. Muller et al. (16 co-authors). 2003. Angiosperm phylogeny based on *matK* sequence information. *Am. J. Bot.* **90**:1758–1776.
- Hiratsuka, J., H. Shimada, R. Whittier et al. (16 co-authors). 1989. The complete sequence of the rice (*Oryza sativa*) chloroplast genome: intermolecular recombination between distinct tRNA genes accounts for a major plastid DNA inversion during the evolution of the cereals. *Mol. Gen. Genet.* **217**:185–194.
- Huelsenbeck, J. P. 2002. Testing a covariotide model of DNA substitution. *Mol. Biol. Evol.* **19**:698–707.
- Hupfer, H., M. Swiatek, S. Hornung, R. G. Herrmann, R. M. Maier, W. L. Chiu, and B. Sears. 2000. Complete nucleotide sequence of the *Oenothera elata* plastid chromosome, representing plastome I of the five distinguishable *Oenothera* plastomes. *Mol. Gen. Genet.* **263**:581–585.
- Jansen, R. K., L. A. Raubeson, J. L. Boore et al. (15 co-authors). 2005. Methods for obtaining and analyzing whole chloroplast genome sequences. *Methods Enzymol.* **395**:348–384.
- Jones, D. T., W. R. Taylor, and J. M. Thornton. 1992. The rapid generation of mutation data matrices from protein sequences. *Comput. Appl. Biosci.* **8**:275–282.
- Judd, W. S., and R. G. Olmstead. 2004. A survey of tricolpate (eudicot) phylogenetic relationships. *Am. J. Bot.* **91**:1627–1644.
- Kato, T., T. Kaneko, S. Sato, Y. Nakamura, and S. Tabata. 2000. Complete structure of the chloroplast genome of a legume, *Lotus japonicus*. *DNA Res.* **7**:323–330.
- Kellogg, E. 2000. Evolutionary history of the grasses. *Plant Physiol.* **125**:1198–1205.
- Kim, K.-J., and H.-L. Lee. 2004. Complete chloroplast genome sequence from korea ginseng (*Panax schinseng* Nees) and comparative analysis of sequence evolution among 17 vascular plants. *DNA Res.* **11**:247–261.
- Kim, S., M.-J. Yoo, V. A. Albert, J. S. Farris, P. S. Soltis, and D. E. Soltis. 2004. Phylogeny and diversification of B-function MADS-box genes in angiosperms: evolutionary and functional implications of a 260-million-year-old duplication. *Am. J. Bot.* **91**:2102–2118.
- Kolaczowski, B., and J. W. Thornton. 2004. Performance of maximum parsimony and likelihood phylogenetics when evolution is heterogeneous. *Nature* **431**:980–984.
- Kramer, E. M., M. A. Jaramillo, and V. S. Di Stilio. 2004. Patterns of gene duplication and functional evolution during the diversification of the AGAMOUS subfamily of MADS box genes in angiosperms. *Genetics* **166**:1011–1023.
- Kugita, M., A. Kaneko, Y. Yamamoto, Y. Takeya, T. Matsumoto, and K. Yoshinaga. 2003a. The complete nucleotide sequence of the hornwort (*Anthoceros formosae*) chloroplast genome:

- insight into the earliest land plants. *Nucleic Acids Res.* **31**:716–721.
- Kugita, M., Y. Yamamoto, T. Fujikawa, T. Matsumoto, and K. Yoshinaga. 2003*b*. RNA editing in hornwort chloroplasts makes more than half the genes functional. *Nucleic Acids Res.* **31**:2417–2423.
- Lake, J. A. 1994. Reconstructing evolutionary trees from DNA and protein sequences: paralinear distances. *Proc. Natl. Acad. Sci. USA* **91**:1455–1459.
- Litt, A., and V. F. Irish. 2003. Duplication and diversification in the APETALA1/FRUITFULL floral homeotic gene lineage: implications for the evolution of floral development. *Genetics* **165**:821–833.
- Lockhart, P. J., C. J. Howe, A. C. Barbrook, A. W. D. Larkum, and D. Penny. 1999. Spectral analysis, systematic bias, and the evolution of chloroplasts. *Mol. Biol. Evol.* **16**:573–576.
- Lockhart, P. J., and D. Penny. 2005. The place of *Amborella* within the radiation of angiosperms. *Trends Plant Sci.* **10**:201–202.
- Lockhart, P. J., M. A. Steel, A. C. Barbrook, D. H. Huson, M. A. Charleston, and C. J. Howe. 1998. A covariotide model explains apparent phylogenetic structure of oxygenic photosynthetic lineages. *Mol. Biol. Evol.* **15**:1183–1188.
- Lockhart, P. J., M. A. Steel, M. D. Hendy, and D. Penny. 1994. Recovering evolutionary trees under a more realistic model of sequence evolution. *Mol. Biol. Evol.* **11**:605–612.
- Lopez, P., D. Casane, and H. Philippe. 2002. Heterotachy, an important process of protein evolution. *Mol. Biol. Evol.* **19**:1–7.
- Lyons-Weiler, J., G. A. Hoelzer, and R. J. Tausch. 1996. Relative apparent synapomorphy analysis (RASA). I: the statistical measurement of phylogenetic signal. *Mol. Biol. Evol.* **13**:749–757.
- Maier, R. M., K. Neckermann, G. L. Igloi, and H. Kossel. 1995. Complete sequence of the maize chloroplast genome: gene content, hotspots of divergence and fine tuning of genetic information by transcript editing. *J. Mol. Biol.* **251**:614–628.
- Martin, W., O. Deusch, N. Stawski, N. Grunheit, and V. Goremykin. 2005. Chloroplast genome phylogenetics: why we need independent approaches to plant molecular evolution. *Trends Plant Sci.* **10**:203–209.
- Mathews, S., and M. J. Donoghue. 1999. The root of angiosperm phylogeny inferred from duplicate phytochrome genes. *Science* **286**:947–950.
- . 2000. Basal angiosperm phylogeny inferred from duplicate phytochromes A and C. *Int. J. Plant Sci.* **161**:S41–S41.
- Ogihara, Y., K. Isono, T. Kojima et al. (18 co-authors). 2000. Chinese spring wheat (*Triticum aestivum* L.) chloroplast genome: complete sequence and contig clones. *Plant Molec. Biol. Rep.* **18**:243–253.
- Ohyama, K., H. Fukuzawa, T. Kohchi et al. 1988. Structure and organization of *Marchantia polymorpha* chloroplast genome. I. Cloning and gene identification. *J. Mol. Biol.* **203**:281–298.
- Parkinson, C. L., K. L. Adams, and J. D. Palmer. 1999. Multigene analyses identify the three earliest lineages of extant flowering plants. *Curr. Biol.* **9**:1485–1488.
- Philippe, H., N. Lartillot, and H. Brinkmann. 2005. Multigene analyses of bilaterian animals corroborate the monophyly of Ecdysozoa, Lophotrochozoa and Protostomia. *Mol. Biol. Evol.* **22**:1246–1253.
- Philippe, H., and J. Laurent. 1998. How good are deep phylogenetic trees? *Curr. Opin. Genet. Dev.* **8**:616–623.
- Phillips, M. J., F. Delsuc, and D. Penny. 2004. Genome-scale phylogeny and the detection of systematic biases. *Mol. Biol. Evol.* **21**:1455–1458.
- Pollock, D. D., D. J. Zwickl, J. A. McGuire, and D. M. Hillis. 2002. Increased taxon sampling is advantageous for phylogenetic inference. *Syst. Biol.* **51**:664–671.
- Qiu, Y. L., J. Lee, F. Bernasconi-Quadroni, D. E. Soltis, P. S. Soltis, M. Zanis, E. A. Zimmer, Z. Chen, V. Savolainen, and M. W. Chase. 1999. The earliest angiosperms: evidence from mitochondrial, plastid and nuclear genomes. *Nature* **402**:404–407.
- . 2000. Phylogeny of basal angiosperms: analyses of five genes from three genomes. *Int. J. Plant Sci.* **161**:S3–S27.
- Rambaut, A., and N. C. Grassly. 1997. Seq-Gen: an application for the Monte Carlo simulation of DNA sequence evolution along phylogenetic trees. *Comput. Appl. Biosci.* **13**:235–238.
- Rokas, A., and P. W. Holland. 2000. Rare genomic changes as a tool for phylogenetics. *Trends Ecol. Evol.* **15**:454–459.
- Sanderson, M. J. 2003. r8s: inferring absolute rates of molecular evolution and divergence times in the absence of a molecular clock. *Bioinformatics* **19**:301–302.
- Sanderson, M. J., J. L. Thorne, N. Wikstrom, and K. Bremer. 2004. Molecular evidence on plant divergence times. *Am. J. Bot.* **91**:1656–1665.
- Sanderson, M. J., M. F. Wojciechowski, J. M. Hu, T. S. Khan, and S. G. Brady. 2000. Error, bias, and long-branch attraction in data for two chloroplast photosystem genes in seed plants. *Mol. Biol. Evol.* **17**:782–797.
- Sato, S., Y. Nakamura, T. Kaneko, E. Asamizu, and S. Tabata. 1999. Complete structure of the chloroplast genome of *Arabidopsis thaliana*. *DNA Res.* **6**:283–290.
- Schmitz-Linneweber, C., R. M. Maier, J. P. Alcaraz, A. Cottet, R. G. Herrmann, and R. Mache. 2001. The plastid chromosome of spinach (*Spinacia oleracea*): complete nucleotide sequence and gene organization. *Plant Mol. Biol.* **45**:307–315.
- Schmitz-Linneweber, C., R. Regel, T. G. Du, H. Hupfer, R. G. Herrmann, and R. M. Maier. 2002. The plastid chromosome of *Atropa belladonna* and its comparison with that of *Nicotiana tabacum*: the role of RNA editing in generating divergence in the process of plant speciation. *Mol. Biol. Evol.* **19**:1602–1612.
- Schneider, H., E. Schuettpelz, K. M. Pryer, R. Cranfill, S. Magallon, and R. Lupia. 2004. Ferns diversified in the shadow of angiosperms. *Nature* **428**:553–557.
- Shimodaira, H., and M. Hasegawa. 1999. Multiple comparisons of log-likelihoods with applications to phylogenetic inference. *Mol. Biol. Evol.* **16**:1114–1116.
- Shinozaki, K., M. Ohme, M. Tanaka et al. (23 co-authors). 1986. The complete nucleotide sequence of tobacco chloroplast genome: its gene organization and expression. *EMBO J.* **5**:2043–2049.
- Simmons, M. P., and M. Miya. 2004. Efficiently resolving the basal clades of a phylogenetic tree using Bayesian and parsimony approaches: a case study using mitogenomic data from 100 higher teleost fishes. *Mol. Phylogenet. Evol.* **31**:351–362.
- Soltis, D. E., V. A. Albert, S. Kim et al. (11 co-authors). 2004*a*. Evolution of the flower. In R. Henry, ed. *Diversity and evolution of plants*. CABI, Wallingford, Oxfordshire.
- Soltis, D. E., V. A. Albert, V. Savolainen et al. (11 co-authors). 2004*b*. Genome-scale data, angiosperm relationships, and “ending incongruence”: a cautionary tale in phylogenetics. *Trends Plant Sci.* **9**:477–483.
- Soltis, D. E., and P. S. Soltis. 2004. *Amborella* not a “basal angiosperm”? Not so fast. *Am. J. Bot.* **91**:997–1001.
- Soltis, P. S., and D. E. Soltis. 2004. The origin and diversification of angiosperms. *Am. J. Bot.* **91**:1614–1626.
- Soltis, P. S., D. E. Soltis, and M. W. Chase. 1999. Angiosperm phylogeny inferred from multiple genes as a tool for comparative biology. *Nature* **402**:402–404.
- Spencer, M., E. Susko, and A. J. Roger. 2005. Likelihood, parsimony, and heterogeneous evolution. *Mol. Biol. Evol.* **22**:1161–1164.

- Steel, M. A. 1994. Recovering a tree from the leaf colorations it generates under a Markov model. *Appl. Math. Lett.* **7**:19–23.
- Stefanović, S., D. W. Rice, and J. D. Palmer. 2004. Long branch attraction, taxon sampling, and the earliest angiosperms: *Amborella* or monocots? *BMC Evol. Biol.* **4**:35.
- Stellari, G. M., M. A. Jaramillo, and E. M. Kramer. 2004. Evolution of the APETALA3 and PISTILLATA lineages of MADS-box-containing genes in the basal angiosperms. *Mol. Biol. Evol.* **21**:506–519.
- Sugiura, C., Y. Kobayashi, S. Aoki, C. Sugita, and M. Sugita. 2003. Complete chloroplast DNA sequence of the moss *Physcomitrella patens*: evidence for the loss and relocation of *rpoA* from the chloroplast to the nucleus. *Nucleic Acids Res.* **31**:5324–5331.
- Swofford, D. L. 2003. PAUP\*: phylogenetic analysis using parsimony (\*and other methods). Version 4. Sinauer Associates, Sunderland, Mass.
- Thollessen, M. 2004. LDDist: a Perl module for calculating LogDet pair-wise distances for protein and nucleotide sequences. *Bioinformatics* **20**:416–418.
- Thompson, J. D., D. G. Higgins, and T. J. Gibson. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**:4673–4680.
- Wakasuki, T., A. Nishikawa, K. Yamada and M. Sugiura. 1998. A complete nucleotide sequence of the plastid genome from a fern, *Psilotum nudum*. *Endocyt. Cell Res.*, **13** (Suppl.):147.
- Wakasugi, T., J. Tsudzuki, S. Ito, K. Nakashima, T. Tsudzuki, and M. Sugiura. 1994. Loss of all *ndh* genes as determined by sequencing the entire chloroplast genome of the black pine *Pinus thunbergii*. *Proc. Natl. Acad. Sci. USA* **91**:9794–9798.
- Williams, J. H., and W. E. Friedman. 2002. Identification of diploid endosperm in an early angiosperm lineage. *Nature* **415**:522–526.
- Wolf, P. G., C. A. Rowe, R. B. Sinclair, and M. Hasebe. 2003. Complete nucleotide sequence of the chloroplast genome from a leptosporangiate fern, *Adiantum capillus-veneris* L. *DNA Res.* **10**:59–65.
- Wyman, S. K., R. K. Jansen, and J. L. Boore. 2004. Automatic annotation of organellar genomes with DOGMA. *Bioinformatics* **20**:3252–3255.
- Yang, Z. 1994. Maximum likelihood phylogenetic estimation from DNA sequences with variable rates over sites: approximate methods. *J. Mol. Evol.* **39**:306–314.
- Zanis, M. J., D. E. Soltis, P. S. Soltis, S. Mathews, and M. J. Donoghue. 2002. The root of the angiosperms revisited. *Proc. Natl. Acad. Sci. USA* **99**:6848–6853.
- Zanis, M. J., P. S. Soltis, Y.-L. Qiu, E. Zimmer, and D. E. Soltis. 2003. Phylogenetic analyses and perianth evolution in basal angiosperms. *Ann. Mo. Bot. Gard.* **90**:129–129.
- Zwickl, D. J., and D. M. Hillis. 2002. Increased taxon sampling greatly reduces phylogenetic error. *Syst. Biol.* **51**:588–598.

William Martin, Associate Editor

Accepted May 27, 2005